

KiaDirect™ Plant Tissue PCR Kit

Cat. No. AD301

Storage: 2X KiaDirect™ PCR kit (+dye) at -20°C for two years; others at room temperature (15-25°C) for two years.

Description

KiaDirect Plant Tissue PCR Kit uses a unique lysis buffer to lyse plant tissues (fresh or frozen). Resulting lysate without purification can be directly used as template for PCR amplification. 2X PCR SuperMix (+dye) is highly resistant to various PCR inhibitors present in plant tissues. PCR product can be directly used for gel electrophoresis. It is recommended to store

Advantages

- Direct amplification from unpurified lysate. Suitable for high-throughput screening.
- Amplification of genomic DNA fragment up to 2 kb

Applications

Plants without high content of polysaccharides or polyphenols

Kit Contents

Component	AD301-01	AD301-02
PDI Buffer	4ml	20ml
PD2 Buffer	4ml	20ml
2X KiaDirect™ PCR Mix (+dye)	1ml	5x1 ml
ddHp	5 ml	25 ml

Please prepare 95°C water bath or heater. If a white precipitate appears, dissolve it by heating at 55°C before use

Genomic DNA Extraction

- Cut 5 mg or 0.5 cm² plant tissues and add it to a tube containing 4 μL of PD1 buffer, vortex.
- Incubate at 95°C for 10 minutes (for plant tissue hard to be lysed, suggest to incubate at 95°C for 30 minutes).
- Add 40 μL of PD2 Buffer and vortex to mix. The lysate can be used as PCR template or stored at 4°C for three months or at -20°C for six months.

Reaction Components

Component	Volume	Final Concentration
Tissue Extract	2-4 μL	as required
Forward Primer (10 μM)	0.4 μL	0.2 μM
Reverse Primer (10 μM)	0.4 μL	0.2 μM
2xKiaDirect™ PCR Mix (+dye)	10 μL	1x
ddHp	Variable	-
Total volume	20 μL	-

Thermal cycling conditions

94°C	5-10 min	} 30-40 cycles
94°C	30 sec	
50-60°C	30 sec	
72°C	1-2 kb/min	
72°C	5-10 min	

FOR RESEARCH USE ONLY

Kiagen Teb Sadra Co.

No 137, Dogol Tower, Oposite of Amie-Atabak, Bagheri Blvd, Motahari Ave,
Tehran, IRAN Tel: 88760179 (5 Lines), 0939-1125991, Fax: 88756622

info@kiagen-biotech.com

www.kiagen-biotech.com

Notes

- Quickly secure the cap of tube for PD1 buffer after use to avoid pH change.
- Avoid repeated freezing and thawing of samples
- Completely thaw the contents in the tube and mix well before use.
- For plant tissues hard to be lysed (e.g. waxy leaves), increase the incubation time at 95°C up to 30 min
- If faint bands are observed, increase the quantity of template used or increase the number of PCR cycles (no more than 40 cycles).
If non-specific amplification bands are observed, adjust the annealing temperature or properly reduce the quantity of template used.
- The extracts can be stored at 4°C for three months or at -20°C for six months.

FOR RESEARCH USE ONLY